

$\mathbb{C}^*$ -extensions of tori, higher Chow groups and applications to incidence equivalence relations for algebraic cycles

by Stefan MÜLLER-STACH, Essen

January 1995

Abstract: Let X be a smooth projective variety over an algebraically closed field k . We repeat Bloch's construction of a $\mathbb{G}_m$ -biextension (torseur) $\mathbb{E}$ over $CH_{\text{hom}}^p(X) \times CH_{\text{hom}}^q(X)$ for $p+q = \dim(X)+1$. First we show that in characteristic zero $\mathbb{E}$ comes via pullback from the Poincaré biextension $\mathbb{P}$ over the corresponding product of intermediate Jacobians - as conjectured by Bloch and Murre. Then the relations between $\mathbb{E}$ and various equivalence relations for algebraic cycles are studied. In particular we reprove Murre's theorem stating that Griffiths' conjecture holds for codimension 2 cycles, i.e. every 2-codimensional cycle which is algebraically and incidence equivalent to zero has torsion Abel-Jacobi invariant.

Key Words: Biextensions, higher Chow groups, algebraic cycles, Abel-Jacobi maps, Deligne-Beilinson cohomology, incidence equivalence, Griffiths' conjecture.

§ 0. Introduction

Let X be a compact Kähler manifold of dimension d . If we fix two integers $0 \leq p, q \leq d$ with $p+q = d+1$, then the two Griffiths intermediate Jacobians $\mathcal{J}^p(X)$ and $\mathcal{J}^q(X)$ are mutually dual. The Poincaré line bundle P on $\mathcal{J}^p(X) \times \mathcal{J}^q(X)$ minus its zero section, denoted by $\mathbb{P}$, is a $\mathbb{C}^*$ -torsor and in fact a biextension in the sense of Grothendieck (SGA 7) and [Mumford 69].

Any two disjoint homologically trivial cycles $Z \in CH_{\text{hom}}^p(X)$, $W \in CH_{\text{hom}}^q(X)$ define an element $\langle Z, W \rangle \in \mathbb{P}$, projecting onto the element $(\psi_p(Z), \psi_q(W)) \in \mathcal{J}^p(X) \times \mathcal{J}^q(X)$, where ψ is the Abel-Jacobi homomorphism.

Bloch has given another construction for a $\mathbb{G}_m$ -torsor $\mathbb{E}$ over $CH_{\text{hom}}^p(X) \times CH_{\text{hom}}^q(X)$ for a projective manifold X over an arbitrary field [Bloch 89]. He also conjectures, that $\mathbb{E}$ may be obtained as the pullback of $\mathbb{P}$ via the Abel-Jacobi map in characteristic zero. We prove this:

Theorem 1. *Let X be a smooth projective variety over $\mathbb{C}$. Then $\mathbb{E}$ is the pullback of $\mathbb{P}$ as torsors under the Abel-Jacobi homomorphism.*

The idea of proof is that the essential point in Bloch's construction factors through Deligne cohomology.

Theorem 1 can be applied to shed some light on the relation between Abel-Jacobi equivalence and incidence equivalence of algebraic cycles. A cycle $W \in CH_{\text{hom}}^q(X)$ is called *incidence equivalent* to zero ($W \sim_{\text{inc}} 0$), if for all pairs (T, B) , with a smooth projective variety T and a cycle $B \in CH^{d+1-q}(T \times X)$, the divisor $B(W) = (pr_T)_*(B \cap pr_X^*(W))$ is linearly equivalent to zero on T (see [Griffiths 69]).

It is easy to show that $\psi(W) = 0$ implies $W \sim_{\text{inc}} 0$. Griffiths conjectured, that the opposite is true modulo torsion if W is assumed to be algebraically equivalent to zero. It was known that this follows from the generalized Hodge conjecture of Grothendieck. Murre ([Murre 85]) has proved the conjecture for codimension two cycles. We show here, that theorem 1 has something to do with this problem. Denote by $\mathbb{E}_W^{\text{alg}}$ the restriction of $\mathbb{E}$ to the fiber over $W \in CH_{\text{alg}}^q(X)$:

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{E}_W^{\text{alg}} \rightarrow CH_{\text{alg}}^p(X) \rightarrow 0$$

We show:

Theorem 2. *Let X be smooth projective over $\mathbb{C}$ and W be algebraically equivalent to zero. Then $\mathbb{E}_W^{\text{alg}}$ is a split extension if and only if $W \sim_{\text{inc}} 0$.*

Here the idea is that incidence equivalence reduces somehow to the situation of points and divisors, which is better understood than the general case. This result brings together the a priori different definitions of incidence equivalence occurring in the literature. Theorem 2 implies a condition for Griffiths' conjecture:

Theorem 3. *Let X be as above and $W \in CH_{\text{alg}}^q(X)$ be such that $\psi(N \cdot W)$ is contained in the dual of $\mathcal{J}_{\text{alg}}^p(X) = \psi(CH_{\text{alg}}^p(X))$ (an abelian subvariety) for some $N \in \mathbb{N}$. Then $W \sim_{\text{inc}} 0$ implies that $\psi(W)$ is torsion in $\mathcal{J}^q(X)$.*

The condition follows again from Grothendieck's generalized Hodge conjecture. But it holds always for codimension two cycles if and therefore we get another proof of Murre's theorem ([Murre85]):

Corollary(Murre's theorem).

Griffiths' conjecture holds for codimension two cycles on a smooth projective manifold X over $\mathbb{C}$, i.e. for every cycle W on X , algebraically and incidence equivalent to zero, $\psi(W)$ is torsion in $\mathcal{J}^2(X)$.

It is easy to see that the proof of Griffiths' conjecture would go through in any codimension p , if the adjoint Λ of the Lefschetz operator were again algebraic. This can be verified for several types of varieties e.g. rational-like varieties and abelian varieties and complete intersections in these. For a special case see [Griffiths-Schmid75]. The paper is organized as follows:

- § 1. The Poincaré biextension.
- § 2. Facts about higher Chow groups.
- § 3. Bloch's construction.
- § 4. The pullback theorem.
- § 5. Abel-Jacobi versus incidence equivalence.

It is a pleasure to thank Jacob Murre for mentioning this problem to me and showing me his unpublished notes ([Murre]) during a visit at Leiden. Furthermore I am grateful to A. Collino, H. Esnault and D. Huybrechts for many helpful remarks and improvements.

§ 1. The Poincaré biextension

Let us first discuss briefly the notion of biextension. We refer to [Mumford 69] and SGA 7 for further details: Let A, B, C be abelian groups. A *biextension* of $B \times C$ by A is a set E on which A acts freely, together with a quotient map $\pi : E \rightarrow B \times C$ and two laws of composition $+_1 : E \times_B E \rightarrow E$, $+_2 : E \times_C E \rightarrow E$ subject to the following conditions:

(1) For all $b \in B$, $\pi^{-1}(b \times C)$ is an abelian group under $+_1$ and the following sequence is exact:

$$0 \rightarrow A \rightarrow \pi^{-1}(b \times C) \rightarrow C \rightarrow 0$$

(2) For all $c \in C$, $\pi^{-1}(B \times c)$ is an abelian group under $+_2$ and the following sequence is exact:

$$0 \rightarrow A \rightarrow \pi^{-1}(B \times c) \rightarrow B \rightarrow 0$$

(3) $+_1, +_2$ are compatible, i.e. for suitable $w, x, y, z \in E$ we have

$$(w +_1 x) +_2 (y +_1 z) = (w +_2 y) +_1 (x +_2 z)$$

There is one prominent example which caused this definition to exist, namely the *Poincaré biextension*:

Let T be a compact complex torus of dimension n . There is a natural line bundle P (the Poincaré line bundle) on $T \times T^\vee$, where $T^\vee = \text{Pic}^0(T)$. As an element of $\text{Pic}(T \times T^\vee)$ it is usually normalized by the two conditions

- (i) $P|_{\{0\} \times T^\vee} \cong \mathcal{O}_{T^\vee}$
- (ii) $P|_{T \times \{\lambda\}}$ represents $\lambda \in \text{Pic}^0(T) = T^\vee$

P is unique under these assumptions. Let $\mathbb{P} = P \setminus \text{zero section}$. Clearly $\mathbb{C}^*$ acts freely on $\mathbb{P}$. $\mathbb{P}$ is called *Poincaré-biextension* associated to T . The projection map $p : \mathbb{P} \rightarrow T \times T^\vee$ makes $\mathbb{P}$ a $\mathbb{C}^*$ -torsor over $T \times T^\vee$. For every $\lambda \in T^\vee$, the inverse image $p^{-1}(T \times \{\lambda\})$ is denoted by $\mathbb{P}_\lambda$ and sits in the exact sequence

$$0 \rightarrow \mathbb{C}^* \rightarrow \mathbb{P}_\lambda \rightarrow T \rightarrow 0.$$

Then $\mathbb{P}_\lambda$ is an extension of abelian groups. It is well known that

$$T^\vee = \text{Pic}^0(T) \cong \text{Ext}^1(T, \mathbb{C}^*)$$

in the category of complex analytic groups. $\mathbb{P}_\lambda$ is exactly the extension of T by $\mathbb{C}^*$ given by λ in this isomorphism. If D is any divisor on T with $c_1(\mathcal{O}(D)) = \lambda$ then we will write also $\mathbb{P}_D$ instead of $\mathbb{P}_\lambda$ and note that $\mathbb{P}_D$ depends only on the class

on D in $\text{Pic}^0(T)$. Now let us consider a special case: Let X be a Kähler manifold of dimension d and p some integer. If we let $T = \mathcal{J}^p(X)$ the p -th intermediate Jacobian of X

$$\mathcal{J}^p(X) = \frac{H^{2p-1}(X, \mathbb{C})}{F^p \oplus H^{2p-1}(X, \mathbb{Z})}$$

then it follows by using Poincaré duality, that $T^\vee$ is given by $\mathcal{J}^q(X)$ where $q = d + 1 - p$. Let

$$\mathbb{P} \rightarrow T \times T^\vee = \mathcal{J}^p(X) \times \mathcal{J}^q(X)$$

be the Poincaré-biextension as defined above. If $W \in CH_{\text{hom}}^q(X)$ with $\lambda = \psi(W) \in \mathcal{J}^q(X)$ and $\mathbb{P}_W := \mathbb{P}_\lambda$ we remark that $\mathbb{P}_W$ depends only on the Abel-Jacobi equivalence class of W , i.e. W is Abel-Jacobi equivalent to zero if and only if $\mathbb{P}_W$ splits as an extension in $\text{Ext}^1(T, \mathbb{C}^*)$.

§ 2. Facts about higher Chow groups

Literature: [Bloch 86], [Murre] for this chapter. Before we repeat Bloch's construction, we recall some properties of *higher Chow groups* as defined by Spencer Bloch.

Let X be a quasi-projective variety over $\mathbb{C}$. Denote by $\Delta^n = \text{Spec}(k[T_0, \dots, T_n] / \langle \sum T_i = 1 \rangle)$ which is isomorphic to affine space $\mathbb{A}_{\mathbb{C}}^n$. There are $n + 1$ natural faces isomorphic to Δ^{n-1} contained in Δ^n , defined by the vanishing of one of the coordinate functions t_i . Let $Z^r(X, n)$ be the free abelian group of cycles $Z \subset X \times \Delta^n$ of codimension r meeting all faces $X \times \Delta^m$ ($m < n$) properly. Then $CH^r(X, n)$ is defined as the n -th homology group of the complex

$$Z^r(X; \cdot) = (\dots Z^r(X, n+1) \xrightarrow{\partial} Z^r(X, n) \xrightarrow{\partial} Z^r(X, n-1) \rightarrow \dots \rightarrow Z^r(X, 0)).$$

The maps ∂ are given by alternating sums of restriction maps to faces. We will need the following properties:

- (1) $CH^*(X, *)$ are covariant (contravariant) functorial for proper (flat) morphisms.
- (2) If $W \subset X$ is closed, we have a long exact sequence
$$\dots CH^*(X, W, n) \rightarrow CH^*(X, n) \rightarrow CH^*(W, n) \rightarrow CH^*(X, W, n-1) \dots$$
- (3) $CH^*(X, 0) = CH^*(X)$ (ordinary Chowgroups).

(4) There is a product for X smooth

$$CH^p(X, q) \otimes CH^r(X, s) \rightarrow CH^{p+r}(X, q + s).$$

(5) There are natural maps to Deligne-Beilinson cohomology, defined in Bloch's article in Contemporary Math. **58** (1986):

$$c_{\mathcal{D}} : CH^p(X, q) \rightarrow H_{\mathcal{D}}^{2p-q}(X, \mathbb{Z}(p)).$$

(6) If X is proper (not necessarily smooth) and $\dim X = d$, then there is a natural *surjective* homomorphism

$$\epsilon : CH^{d+1}(X, 1) \rightarrow \mathbb{C}^*$$

factoring through Deligne-Beilinson cohomology

$$\epsilon : CH^{d+1}(X, 1) \xrightarrow{c_{\mathcal{D}}} H_{\mathcal{D}}^{2d+1}(X, \mathbb{Z}(d+1)) \rightarrow \mathbb{C}^*$$

obtained as follows: Let $\pi : X \rightarrow \text{Spec}(\mathbb{C})$ be the natural morphism. Then ϵ is given by the direct image map $\pi_* : CH^{d+1}(X, 1) \rightarrow CH^1(\text{Spec}(\mathbb{C}), 1)$, since it is a straightforward exercise to show that

$$CH^1(\text{Spec}(\mathbb{C}), 1) \cong H_{\mathcal{D}}^1(\text{Spec}(\mathbb{C}), \mathbb{Z}(1)) \cong \mathbb{C}/\mathbb{Z}(1)$$

via the classes in (5) and the latter group can be identified with $\mathbb{C}^*$. We claim:

(1) ϵ is surjective.

(2) It factors through Deligne-Beilinson cohomology.

Proof for (1) and (2)

(1): By definition $CH^1(\text{Spec}(\mathbb{C}), 1) = Z^1(\text{Spec}(\mathbb{C}), 1)/\text{Im}(\partial)$, since $Z^1(\text{Spec}(\mathbb{C}), 0) = 0$. In the same way $CH^{d+1}(X, 1) = Z^{d+1}(X, 1)/\text{Im}(\partial)$. Hence in both groups the elements are represented by finite sums of points. It is clear that the map $Z^{d+1}(X, 1) \rightarrow Z^1(\text{Spec}(\mathbb{C}), 1)$ induced by π_* is surjective. Hence we deduce (1).

(2): By the functorial properties of the Deligne-Beilinson classes we get a commutative diagram:

$$\begin{array}{ccc} CH^{d+1}(X, 1) & \xrightarrow{\pi_*} & CH^1(\text{Spec}(\mathbb{C}), 1) \\ \downarrow c_{\mathcal{D}} & & \downarrow c_{\mathcal{D}} \\ H_{\mathcal{D}}^{2d+1}(X, \mathbb{Z}(d+1)) & \xrightarrow{\pi_*} & H_{\mathcal{D}}^1(\text{Spec}(\mathbb{C}), \mathbb{Z}(1)) \end{array}$$

The right vertical arrow being the identity we get (2). □

§ 3. Bloch's Construction

For this chapter we refer to [Bloch 89].

Let us describe Bloch's construction in detail: Let X be smooth and projective over $\mathbb{C}$ with $\dim X = d$, and $W \in Z_{m-1}(X) = Z^{d+1-m}(X)$ a cycle of dimension $m-1$, which is homologous to zero, denoted by $W \sim_{\text{hom}} 0$ as usual. We will construct an extension

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{E}_W \rightarrow CH_{\text{hom}}^m(X) \rightarrow 0$$

such that they can be glued together to give

$$\mathbb{E} = \bigcup \mathbb{E}_W \rightarrow CH_{\text{hom}}^m(X) \times CH_{\text{hom}}^{d+1-m}(X)$$

a biextension in the sense of § 1. To do this, Bloch first considers the map

$$\Theta_W = \epsilon \circ (\cap W) : CH^m(X, 1) \xrightarrow{\cap W} CH^{d+1}(X, 1) \xrightarrow{\epsilon} \mathbb{C}^*$$

using properties (4) and (6) of higher Chow groups.

Lemma 1. *If $W \sim_{\text{hom}} 0$, then $\Theta_W \equiv 1$.*

Proof. Recall the following fact from Deligne-Beilinson cohomology ([Esnault-Viehweg 88]):

$\mathcal{J}^* \subset H_{\mathcal{D}}^*(X, \mathbb{Z}(*))$ is a square zero ideal, where

$$\mathcal{J}^* = \text{Ker}(H_{\mathcal{D}}^*(X, \mathbb{Z}(*)) \rightarrow H_{\text{Betti}}^*(X, \mathbb{Z}(*)))$$

Now look at the diagram

$$\begin{array}{ccccccc} CH^m(X, 1) & & \xrightarrow{\cap W} & & CH^{d+1}(X, 1) & & \xrightarrow{\epsilon} \mathbb{C}^* \\ \downarrow c_{\mathcal{D}} & & & & \downarrow c_{\mathcal{D}} & & \parallel \\ H_{\mathcal{D}}^{2m-1}(X, \mathbb{Z}(m)) & & \xrightarrow{\cap c_{\mathcal{D}}(W)} & & H_{\mathcal{D}}^{2d+1}(X, \mathbb{Z}(d+1)) & & \xrightarrow{\epsilon} \mathbb{C}^* \end{array}$$

But $W \sim_{\text{hom}} 0$ implies $c_{\mathcal{D}}(W) \in \mathcal{J}^*(X)$, and also $H_{\mathcal{D}}^{2m-1}(X, \mathbb{Z}(m)) \subset \mathcal{J}^*(X)$. Hence the lemma follows. $\square$

To construct $\mathbb{E}_W$ consider the long exact sequence for the pair $(X, |W|)$:

$$\dots \rightarrow CH^m(X, 1) \rightarrow CH^m(|W|, 1) \rightarrow CH^m(X, |W|, 0) \rightarrow CH^m(X, 0) \rightarrow 0$$

where $|W| = \text{supp}(W)$ as usual. Since $|W|$ is proper of dimension $m-1$, we have a surjective map $\epsilon : CH^m(|W|, 1) \rightarrow \mathbb{C}^*$ by property (6). But $\Theta_W \equiv 1$, hence ϵ factors through

$$A := \frac{CH^m(|W|, 1)}{\text{Im } CH^m(X, 1)} \rightarrow \mathbb{C}^*.$$

Consider the commutative diagram

$$\begin{array}{ccccccc}
0 & \rightarrow & A & \xrightarrow{i} & CH^m(X, |W|, 0) & \rightarrow & CH^m(X) \rightarrow 0 \\
& & \downarrow \epsilon & & \downarrow & & \parallel \\
1 & \rightarrow & \mathbb{C}^* & \rightarrow & \mathbb{E}_W & \rightarrow & CH^m(X) \rightarrow 0
\end{array}$$

where we have defined $\mathbb{E}_W$ as

$$CH^m(X, |W|, 0)/i(\text{Ker}(\epsilon))$$

as the pushout of the upper line via ϵ . By abuse of language, we also denote by $\mathbb{E}_W$ the restriction

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{E}_W \rightarrow CH_{\text{hom}}^m(X) \rightarrow 0$$

to $CH_{\text{hom}}^m(X)$. Bloch proves several results for $\mathbb{E}_W$:

If W is rationally equivalent to W' , then $\mathbb{E}_W$ is canonically isomorphic to $\mathbb{E}_{W'}$. In particular, if $W \sim_{\text{rat}} 0$, then $\mathbb{E}_W$ splits as an extension of groups. We prove a little bit more in § 4.

The $\mathbb{E}_W$ may be glued together, to obtain a $\mathbb{C}^*$ -torseur (biextension)

$$\mathbb{E} \rightarrow CH_{\text{hom}}^m(X) \times CH_{\text{hom}}^{d+1-m}(X)$$

satisfying the axioms of § 1.

We refer the reader to [Bloch 89] for this construction, since we don't use the full torseur $\mathbb{E}$ for our applications. In fact by our theorem 1 this follows from the statement for $\mathbb{P}$.

§ 4. The Pullback Theorem

Here we prove the main result of this paper. Let X be a smooth projective variety over $\mathbb{C}$.

Lemma 2. *If $W \sim_{AJ} 0$ (Abel-Jacobi equivalent to zero), then $\mathbb{E}_W$ splits.*

Proof. We use the functorial Deligne-Beilinson classes $c_{\mathcal{D}}$ from (5), §2. Consider the commutative diagram

$$\begin{array}{ccccccc}
A & \rightarrow & CH_{\text{hom}}^m(X, |W|, 0) & \rightarrow & CH_{\text{hom}}^m(X) & \rightarrow & 0 \\
\downarrow c_{\mathcal{D}} & & \downarrow & & \downarrow & \text{Abel - Jacobi} & \\
H_{\mathcal{D}}^{2m-1}(|W|; \mathbb{Z}(m)) & \rightarrow & H_{\mathcal{D}}^{2m}(X, |W|; \mathbb{Z}(m))_{\text{hom}} & \rightarrow & \mathcal{J}^m(X) & \rightarrow & 0
\end{array}$$

$\epsilon : A \rightarrow \mathbb{C}^*$ is surjective and by §2.(6) factors through

$$H_{\mathcal{D}}^{2m-1}(|W|; \mathbb{Z}(m)) \rightarrow \bigoplus \mathbb{C}^* \xrightarrow{\Sigma} \mathbb{C}^*$$

by summation over all irreducible components of $|W|$. Define $K := \text{Ker}(A \rightarrow \mathbb{C}^*)$ and recalling that $\mathbb{E}_W = CH_{\text{hom}}^m(X, |W|, 0)/i(K)$ we obtain the diagram

$$\begin{array}{ccccccc} & & T & & T' & & \\ & & \downarrow & & \downarrow & & \\ 0 & \rightarrow & A/K & \rightarrow & \mathbb{E}_W & \rightarrow & CH_{\text{hom}}^m(X) \rightarrow 0 \\ & & \downarrow \epsilon & & \downarrow & & \downarrow \\ 1 & \rightarrow & \mathbb{C}^* & \rightarrow & \mathbb{P}_W & \rightarrow & \mathcal{J}^m(X) \rightarrow 0 \end{array} \quad \text{Abel - Jacobi}$$

where $\mathbb{P}_W$ is as in § 1, $T = \text{Ker}(\mathbb{E}_W \rightarrow \mathbb{P}_W)$ and $T' = \text{Ker}(CH_{\text{hom}}^m(X) \rightarrow \mathcal{J}^m(X))$. Since ϵ is an isomorphism, $T \cong T'$ by the snake lemma. It follows that $\mathbb{E}_W$ is - as an extension - the pullback of $\mathbb{P}_W$ via the Abel-Jacobi map and therefore is split if W is Abel-Jacobi equivalent to zero. This proves the lemma. $\square$

As a corollary we get:

Theorem 1. $\mathbb{E}_W$ is the pullback of $\mathbb{P}_W$ as extensions via the Abel-Jacobi map. In particular $\mathbb{E}$ is the pullback of $\mathbb{P}$ as torseurs via the Abel-Jacobi map.

§ 5. Abel-Jacobi versus Incidence Equivalence

The notion of “incidence equivalence” was introduced by Griffiths ([Griffiths 69]). Let X be projective, smooth of dimension d over $\mathbb{C}$.

Definition: A cycle $W \in CH_{\text{hom}}^i(X)$ is called *incidence equivalent to zero*, i.e. $W \sim_{\text{inc}} 0$, if for all pairs (T, B) with T a smooth, projective variety and $B \in CH^{d+1-i}(T \times X)$,

$$B(W) := (pr_T)_*(B \cap pr_X^* W) = \{t \in T \mid B_t \cap W \neq \emptyset\}$$

is linearly equivalent to zero in $\text{Pic}^0(T)$.

Remark: $B(W)$ is a divisor by a dimension count and in $\text{Pic}^0(T)$, since $W \sim_{\text{hom}} 0$. If $i = 1$ we get back the notion of rational equivalence for divisors (= Abel-Jacobi equivalence).

Bloch defines $W \sim_{\text{inc}} 0$ if $\mathbb{E}_W$ is a split extension. However it is not a priori clear that this gives the same definition. See the discussion below.

Lemma 3. *If $W \sim_{A\mathcal{J}} 0$, then also $W \sim_{\text{inc}} 0$.*

Proof. B is a correspondence

$$CH_{\text{hom}}^i(X) \rightarrow CH_{\text{hom}}^1(T) = \text{Pic}^0(T)$$

via $W \mapsto B(W)$. Consider the diagram (ψ = Abel-Jacobi homomorphism)

$$\begin{array}{ccc} CH_{\text{hom}}^i(X) & \xrightarrow{B} & CH_{\text{hom}}^1(T) = \text{Pic}^0(T) \\ \downarrow \psi & & \downarrow S \\ \mathcal{J}^i(X) & \xrightarrow{\mathcal{J}B} & \mathcal{J}^1(T) \end{array}$$

The claim follows. $\square$

Remark: Griffiths has conjectured, that the opposite is true modulo torsion for cycles algebraically equivalent to zero: If $W \sim_{\text{inc}} 0$, then for some $N \in \mathbb{N}$: $N \cdot W \sim_{A\mathcal{J}} 0$. This is true for $i = 1, d$ without torsion and for $i = 2$ by [Murre 85]. In any case it follows from the generalized Hodge conjecture of Grothendieck, as we will see below.

In the remaining discussion let $i = d + 1 - m$ and W a cycle of dimension $m - 1$, hence codimension $d + 1 - m$ which is homologous to zero.

We want to investigate the relation between $W \sim_{\text{inc}} 0$ and the splitting of $\mathbb{E}_W$ in the sense of [Bloch89]. To do this consider the action of the correspondence B on the higher Chow groups. $B(W) \in \text{Pic}^0(T)$ gives rise to the exact sequence

$$A_{B(W)} = \frac{CH^t(|B(W)|, 1)}{\text{Im } CH^t(T, 1)} \rightarrow CH^t(T, |B(W)|, 0) \rightarrow CH^t(T) \rightarrow 0$$

where $t = \dim T$ and the corresponding extension $\mathbb{E}_{B(W)}$ is sitting in

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{E}_{B(W)} \rightarrow CH_{\text{hom}}^t(T) \rightarrow 0$$

and is pulled back via the Albanese map from $\mathbb{P}_W$:

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{P}_{B(W)} \rightarrow \text{Alb}(T) \rightarrow 0.$$

The correspondence B induces maps $B^\sharp = B^{-1} : CH_{\text{hom}}^t(T) \rightarrow CH_{\text{hom}}^m(X)$ then giving rise to a commutative diagram

$$\begin{array}{ccccccc} 0 & \rightarrow & A_{B(W)} & \rightarrow & CH^t(T, |B(W)|, 0) & \rightarrow & CH^t(T) \rightarrow 0 \\ & & \downarrow B^\sharp & & \downarrow B^\sharp & & \downarrow B^\sharp \\ 0 & \rightarrow & A_W & \rightarrow & CH^m(X, |W|, 0) & \rightarrow & CH^m(X) \rightarrow 0 \end{array}$$

This can be seen as follows: By Chow's moving lemma, cycles in $CH^m(X)$ and $CH^m(X, |W|)$ may be assumed to have support disjoint from W . The same holds for T and $B(W)$. Now if a zero cycle on T does not meet $B(W)$ then $B^\sharp(Z)$ will

not meet W . This explains the maps in the diagram. Everything commutes by the functorial properties of § 2.

We thus obtain a diagram

$$\begin{array}{ccccccc} 1 & \rightarrow & \mathbb{C}^* & \rightarrow & \mathbb{E}_{B(W)} & \rightarrow & CH_{\text{hom}}^t(T) \rightarrow 0 \\ & & || & & \downarrow B^\sharp & & \downarrow B^\sharp \\ 1 & \rightarrow & \mathbb{C}^* & \rightarrow & \mathbb{E}_W & \rightarrow & CH_{\text{hom}}^m(X) \rightarrow 0 \end{array}$$

inducing the identity on $\mathbb{C}^*$, where we assumed that B is reduced without loss of generality.

This is the main input in order to prove:

Lemma 4. (a) If $\mathbb{E}_W$ splits then also $\mathbb{E}_{B(W)}$ splits for all pairs (T, B) .

(b) Assume additionally that $W \sim_{\text{alg}} 0$. Then:

$\mathbb{E}_{B(W)}$ splits for all pairs $(T, B) \iff \mathbb{E}_W^{\text{alg}}$ splits, where $\mathbb{E}_W^{\text{alg}}$ is the subextension

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{E}_W^{\text{alg}} \rightarrow CH_{\text{alg}}^m(X) \rightarrow 0$$

Proof. (a) follows directly from the commutative diagram for every pair (T, B) . To prove (b) suppose $W \sim_{\text{alg}} 0$ and that $\mathbb{E}_{B(W)}$ splits for all pairs (T, B) . We have to show that $\mathbb{E}_W^{\text{alg}}$ splits. Take (T, B) such that

$$B^\sharp : \text{Alb}(T) \rightarrow \mathcal{J}^m(X)$$

parametrizes the whole image $\mathcal{J}_{\text{alg}}^m(X)$ of the Abel-Jacobi map for cycles algebraically equivalent to zero. T can be chosen to be an abelian variety itself such that $B^\sharp$ becomes an isogeny. It follows that the subextension

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{P}_W^{\text{alg}} \rightarrow \mathcal{J}_{\text{alg}}^m(X) \rightarrow 0$$

of $\mathbb{P}$ splits by the Deligne cohomology version of the commutative diagram above. Now by theorem 1 and its proof, the extension class of $\mathbb{E}_W^{\text{alg}}$ is pulled back from $\text{Ext}^1(\psi(CH_{\text{alg}}^m(X)), \mathbb{C}^*) = \text{Ext}^1(\mathcal{J}_{\text{alg}}^m(X), \mathbb{C}^*)$. (b) follows. $\square$

As a corollary we get:

Theorem 2. Assume $W \sim_{\text{alg}} 0$. Then $W \sim_{\text{inc}} 0$ if and only if $\mathbb{E}_W^{\text{alg}}$ splits.

Proof. $W \sim_{\text{inc}} 0$ iff $B(W)$ is linearly equivalent to zero by definition, hence iff $\mathbb{P}_{B(W)}$ splits. But for divisors the splittings of $\mathbb{E}$ and $\mathbb{P}$ are equivalent (as one can see by restricting to suitable curve) and hence this holds if and only if $\mathbb{E}_{B(W)}$ splits. Hence by lemma 4 (b) we are finished. $\square$

It remains to discuss *Griffiths' conjecture*: Let $W \sim_{\text{alg}} 0$. If $W \sim_{\text{inc}} 0$ then some multiple of W is Abel-Jacobi equivalent to zero. With the help of theorem 2 it is sufficient to show: If $\mathbb{E}_W$ splits, then for some $N \in \mathbb{N}$: $\mathbb{P}_{N \cdot W}$ splits.

Lemma 5. *Suppose the generalized Hodge conjecture (GHC) holds for $\mathcal{J}^m(X)$, i.e. the largest abelian subvariety $\mathcal{J}_{\text{ab}}^m(X)$ of $\mathcal{J}^m(X)$ is parametrized by algebraic cycles. Then for $W \in CH_{m-1}(X)$, $W \sim_{\text{alg}} 0$, we have: If $\mathbb{E}_W$ splits, then for some $N \in \mathbb{N}$, $\mathbb{P}_{N \cdot W}$ splits.*

Proof. Remark that $\mathcal{J}_{\text{ab}}^m(X)$ and $\mathcal{J}_{\text{ab}}^{d+1-m}(X)$ are always dual to each other modulo isogeny, since the dual of a Hodge substructure is again one. Therefore GHC implies that $\mathcal{J}_{\text{alg}}^m(X) = \mathcal{J}_{\text{ab}}^m(X)$ and that $\psi(N \cdot W)$ lies in the dual abelian variety of $\mathcal{J}_{\text{alg}}^m(X) = \psi(CH_{\text{alg}}^m(X))$ for some $N \in \mathbb{N}$. Hence it is sufficient to show that the extension

$$1 \rightarrow \mathbb{C}^* \rightarrow \mathbb{P}_{N \cdot W, \text{alg}} \rightarrow \mathcal{J}_{\text{alg}}^m(X) \rightarrow 0$$

splits. This in turn is implied by the splitting of $\mathbb{E}_W$ and hence of $\mathbb{E}_{N \cdot W}^{\text{alg}}$ by the proof of lemma 4. $\square$

Actually we have proved more:

Theorem 3. *Let $W \in CH_{m-1}(X)$ be algebraically equivalent to zero. Then if $\psi(N \cdot W)$ is contained in the dual of $\mathcal{J}_{\text{alg}}^m(X)$, the splitting of $\mathbb{E}_W$ implies the splitting of $\mathbb{P}_{N \cdot W}$.*

A little generalization of the argument in Lemma 5.2. of [Murre85] gives another proof of Murre's theorem:

Corollary(Murre's theorem).

Griffiths' conjecture holds for codimension two cycles on a smooth projective manifold X over $\mathbb{C}$, i.e. for every cycle W on X , algebraically and incidence equivalent to zero, $\psi(W)$ is torsion in $\mathcal{J}^2(X)$.

Proof. We have to verify the assumptions of theorem 3. Let T be the universal cover (tangent space) of $\mathcal{J}_{\text{alg}}^2(X)$ and $H_a^3(X) = T \oplus \bar{T}$. This is a sub Hodge structure of $H^3(X, \mathbb{C})$ contained in $H^{1,2}(X) \oplus H^{2,1}(X)$. Now $\psi(W)$ is contained in the dual of $\mathcal{J}_{\text{alg}}^{d-1}(X)$ ($d = \dim(X)$) if we can show that the cup product

$$H_a^3(X) \otimes H_a^{2d-3}(X) \rightarrow \mathbb{C}$$

has no left kernel. Let L be the Lefschetz operator, i.e. cup product with the hyperplane class. It is algebraic and hence maps $H_a^k(X)$ to $H_a^{k+2}(X)$.

Let us repeat the proof of Murre's Lemma 5.2. for convenience. Assume there is a nonzero element $v \in H_a^3(X)$ in the left kernel and decompose it as $v = v_0 + Lv_1$ according to Lefschetz decomposition. The v_k are primitive. In particular $v_1 \in H_a^1(X)$ which is the whole of $H^1(X, \mathbb{C})$. Hence Lv_1 and v_0 are in $H_a^3(X)$. Now decompose $v_0 = w_1 + w_2$ into types $(1, 2)$ and $(2, 1)$. Since $H_a^{2p-1}(X)$ is a sub Hodge structure, w_1 and w_2 and their complex conjugates are again in $H_a^{2p-1}(X)$. By symmetry we may assume $w_1 \neq 0$ unless $v_0 = 0$. In the first case set $z := L^{d-3}(\bar{w}_1) \in H_a^{2d-3}(X)$, otherwise decompose $v_1 = y_1 + y_2$ into types with $y_1 \neq 0$ (by symmetry) and set $z := L^{d-2}(\bar{y}_1)$. Then for type reasons in the first case $v \cup z = v_0 \cup z = L^{d-3}(w_1 \cup \bar{w}_1) \neq 0$ by primitivity of w_1 , a contradiction. In the second case $v \cup z = Lv_1 \cup z = L^{d-1}(y_1 \cup \bar{y}_1) \neq 0$ by primitivity of y_1 , a contradiction. $\square$

Remark: It is easy to see that the proof of Griffiths' conjecture would go through in any codimension p , if in the Lefschetz decomposition of $v \in H_a^{2p-1}(X)$ every term were again in $H_a^{2p-2k-1}(X)$. This can be verified for several types of varieties e.g. rational-like varieties and abelian varieties and complete intersections in these. For a special case see [Griffiths-Schmid75].

References

- [Bloch 86] S. Bloch: Algebraic cycles and higher K -theory. Adv. in Math. 61, 167 - 304 (1986)
- [Bloch 89] S. Bloch: Cycles and Biextensions, in Cont. Math. 83, 19 - 30 (1989)
- [Carlson 85] J. Carlson: The geometry of the extension class of a mixed Hodge structure, in Proc. Bowdoin conference of the AMS Vol.2, 199-222 (1987)
- [Esnault-Viehweg 88] H. Esnault, E. Viehweg: Deligne-Beilinson Cohomology, Perspectives in Mathematics, Vol. 4, Acad. Press, 43-91 (1988)
- [Griffiths 69] Ph. Griffiths: Some results on algebraic cycles on algebraic manifolds, Proc. Bombay conference, 93 - 191 (1969)
- [Griffiths-Schmid75] Ph. Griffiths, W. Schmid: Recent developments in Hodge theory, in Discrete subgroups of Liegroups, Bombay Coll. 1975, Oxford Univ. Press, 31-127 (1975)
- [Hain 90] R. Hain: Biextensions and heights associated to curves of odd genus; Duke Math. J. 61, 859 - 898 (1990)
- [Mumford 69] D. Mumford: Biextensions of formal groups, Proc. Bombay Colloquium 1968, Oxford Univ. Press, 307-322 (1969)
- [Murre] unpublished notes
- [Murre 85] J. Murre: Abel Jacobi equivalence versus incidence equivalence for algebraic cycles of codimension two, Topology 24 (3), 361 - 367 (1985)